

Package: rgsp (via r-universe)

September 2, 2024

Type Package

Title Repetitive Group Sampling Plan Based on Cpk

Version 0.2.0

Author Muhammad Yaseen [aut, cre], Muhammad Aslam [aut, ctb], Sami Ullah [aut, ctb], Muhammad Kashif [aut, ctb]

Maintainer Muhammad Yaseen <myaseen208@gmail.com>

Description Functions to calculate Sample Number and Average Sample Number for Repetitive Group Sampling Plan Based on Cpk as given in Aslam et al. (2012) (<DOI:10.1080/00949655.2012.663374>).

Depends R (>= 3.1)

Imports dplyr, magrittr, tibble

License GPL-2

URL <https://github.com/myaseen208/rgsp>,
<https://myaseen208.github.io/rgsp/>

Encoding UTF-8

LazyData true

RoxygenNote 6.1.0

Note Department of Mathematics and Statistics, University of Agriculture Faisalabad, Faisalabad-Pakistan.

Suggests testthat

Repository <https://myaseen208.r-universe.dev>

RemoteUrl <https://github.com/myaseen208/rgsp>

RemoteRef HEAD

RemoteSha 054efcccebc5091cb21f61871e090126fb3197da

Contents

rgsp	2
rgsp_asym1	2
rgsp_asym2	3
rgsp_sym	5

Index**7**

rgsp

*Repetitive Group Sampling Plan Based on Cpk***Description**

The rgsp package provides functionalities to calculate Sample Number and Average Sample Number for a Repetitive Group Sampling Plan based on Cpk as given in Aslam et al. (2013).

Author(s)

1. Muhammad Yaseen (<myaseen208@gmail.com>)
2. Muhammad Aslam (<aslam_ravian@hotmail.com>)
3. Sami Ullah (<samiullahuos@gmail.com>)
4. Muhammad Kashif (<mkashif@uaf.edu.pk>)

References

Aslam, M., Wu, C., Jun, C., Azam, M. and Itay, N. (2013). Developing a variables repetitive group sampling plan based on process capability index Cpk with unknown mean and variance. *Journal of Statistical Computation and Simulation*. **83**(8):1507-1517. (<https://www.tandfonline.com/doi/abs/10.1080/00949655.2012.6>)

rgsp_asym1

*Repetitive Group Sampling Plan Based on Cpk under asymmetric Case 1***Description**

Calculates Sample Number and Average Sample Number for Repetitive Group Sampling Plan based on Cpk under asymmetric case 1 as given in Aslam et al. (2013)

Usage

```
## Default S3 method:
rgsp_asym1(.p1, .p2, .alpha, .beta, .nums, .rep)
```

Arguments

.p1	Acceptable Quality Level (AQL) Probability
.p2	Limiting Quality Level (LQL) Probability
.alpha	Producer's alpha-risk
.beta	Consumer's beta-risk
.nums	Number of samples with replacement at each iteration
.rep	Number of iterations

Value

Sample Number and Average Sample Number

Author(s)

1. Muhammad Yaseen (<myaseen208@gmail.com>)
2. Muhammad Aslam (<aslam_ravian@hotmail.com>)
3. Sami Ullah (<samiullahuos@gmail.com>)
4. Muhammad Kashif (<mkashif@uaf.edu.pk>)

References

Aslam, M., Wu, C., Jun, C., Azam, M. and Itay, N. (2013). Developing a variables repetitive group sampling plan based on process capability index Cpk with unknown mean and variance. *Journal of Statistical Computation and Simulation*. **83**(8):1507-1517. (<https://www.tandfonline.com/doi/abs/10.1080/00949655.2012.6>)

Examples

```
rgsp_asym1(
  .p1 = 0.001
  , .p2 = 0.003
  , .alpha = 0.050
  , .beta = 0.100
  , .nums = 10000
  , .rep = 10 # 1000
)
```

rgsp_asym2

Repetitive Group Sampling Plan Based on Cpk under asymmetric Case 2

Description

Calculates Sample Number and Average Sample Number for Repetitive Group Sampling Plan based on Cpk under asymmetric case 2 as given in Aslam et al. (2013)

Usage

```
## Default S3 method:
rgsp_asym2(.p1, .p2, .alpha, .beta, .nums, .rep)
```

Arguments

.p1	Acceptable Quality Level (AQL) Probability
.p2	Limiting Quality Level (LQL) Probability
.alpha	Producer's alpha-risk
.beta	Consumer's beta-risk
.nums	Number of samples with replacement at each iteration
.rep	Number of iterations

Value

Sample Number and Average Sample Number

Author(s)

1. Muhammad Yaseen (<myaseen208@gmail.com>)
2. Muhammad Aslam (<aslam_ravian@hotmail.com>)
3. Sami Ullah (<samiullahuos@gmail.com>)
4. Muhammad Kashif (<mkashif@uaf.edu.pk>)

References

Aslam, M., Wu, C., Jun, C., Azam, M. and Itay, N. (2013). Developing a variables repetitive group sampling plan based on process capability index Cpk with unknown mean and variance. *Journal of Statistical Computation and Simulation*. **83**(8):1507-1517. (<https://www.tandfonline.com/doi/abs/10.1080/00949655.2012.6>)

Examples

```
rgsp_asym2(  
  .p1      = 0.001  
  , .p2    = 0.003  
  , .alpha = 0.050  
  , .beta  = 0.100  
  , .nums  = 10000  
  , .rep   = 10 # 1000  
)
```

rgsp_sym

Repetitive Group Sampling Plan Based on Cpk under Symmetric Case

Description

Calculates Sample Number and Average Sample Number for Repetitive Group Sampling Plan based on Cpk under symmetric case as given in Aslam et al. (2013)

Usage

```
## Default S3 method:  
rgsp_sym(.p1, .p2, .alpha, .beta, .nums, .rep)
```

Arguments

.p1	Acceptable Quality Level (AQL) Probability
.p2	Limiting Quality Level (LQL) Probability
.alpha	Producer's alpha-risk
.beta	Consumer's beta-risk
.nums	Number of samples with replacement at each iteration
.rep	Number of iterations

Value

Sample Number and Average Sample Number

Author(s)

1. Muhammad Yaseen (<myaseen208@gmail.com>)
2. Muhammad Aslam (<aslam_ravian@hotmail.com>)
3. Sami Ullah (<samiullahuos@gmail.com>)
4. Muhammad Kashif (<mkashif@uaf.edu.pk>)

References

Aslam, M., Wu, C., Jun, C., Azam, M. and Itay, N. (2013). Developing a variables repetitive group sampling plan based on process capability index Cpk with unknown mean and variance. *Journal of Statistical Computation and Simulation*. **83**(8):1507-1517. (<https://www.tandfonline.com/doi/abs/10.1080/00949655.2012.6>)

Examples

```
rgsp_sym(  
  .p1 = 0.0010  
  , .p2 = 0.0020  
  , .alpha = 0.0500  
  , .beta = 0.1000  
  , .nums = 10000  
  , .rep = 10 # 1000  
)
```

Index

rgsp, [2](#)
rgsp-package (rgsp), [2](#)
rgsp_asym1, [2](#)
rgsp_asym2, [3](#)
rgsp_sym, [5](#)